

APPROXIMATIONS OF INJECTIVE MODULES AND FINITISTIC DIMENSION

F. HUARD AND D. SMITH

ABSTRACT. Let Λ be an artin algebra and let $\mathcal{P}_\Lambda^{<\infty}$ the category of finitely generated right Λ -modules of finite projective dimension. We show that $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite in $\text{mod } \Lambda$ if and only if the direct sum E of the indecomposable Ext-injective modules in $\mathcal{P}_\Lambda^{<\infty}$ form a tilting module in $\text{mod } \Lambda$. Moreover, we show that in this case E coincides with the direct sum of the minimal right $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the indecomposable Λ -injective modules and that the projective dimension of E equal to the finitistic dimension of Λ .

Given an artin algebra Λ , we let $\mathcal{P}_\Lambda^{<\infty}$ denote the class of all finitely generated right Λ -modules of finite projective dimension. The finitistic dimension of Λ , $\text{fin.dim. } \Lambda$, is then defined to be the supremum of the projective dimensions of the modules in $\mathcal{P}_\Lambda^{<\infty}$.

It is still an open question whether $\text{fin.dim. } \Lambda$ is always finite. However, when $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite, it was shown by Auslander and Reiten that $\text{fin.dim. } \Lambda$ is finite. They showed in [2] that under this hypothesis, the finitistic dimension of Λ is equal to the maximum of the projective dimensions of the minimal right $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the simple Λ -modules. In this note, we show that this result remains valid when simples are replaced by indecomposable injectives. Moreover, and it is crucial to our proof, the direct sum of these $\mathcal{P}_\Lambda^{<\infty}$ -approximations of indecomposable injectives turns out to be a tilting module. This is not true in general for the $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the simple Λ -modules.

Conversely, by using results from [1, 3], we show that if the direct sum E of the indecomposable Ext-injective modules in $\mathcal{P}_\Lambda^{<\infty}$ form a tilting module, then $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite and E coincides with the direct sum of the minimal right $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the indecomposable Λ -injective modules. This complements the results of [4] that were obtained for contravariantly finite and resolving subcategories of $\mathcal{P}_\Lambda^{\leq n}$, that is the category of all finitely generated right Λ -modules of projective dimension at most n , with $n \in \mathbb{N}$.

Let $\text{Mod } \Lambda$ denote the category of all right Λ -modules and $\text{mod } \Lambda$ be the category of finitely generated right Λ -modules. Let $\mathcal{X}$ be a subcategory of $\text{mod } \Lambda$. By a *right $\mathcal{X}$ -approximation* of a module $M \in \text{mod } \Lambda$, we mean a morphism $f_M : X_M \rightarrow M$, with $X_M \in \mathcal{X}$, such that any morphism $f : X \rightarrow M$, with $X \in \mathcal{X}$, factors through f_M . We say that $\mathcal{X}$ is *contravariantly finite* if each $M \in \text{mod } \Lambda$ admits a right $\mathcal{X}$ -approximation.

A module $X \in \mathcal{X}$ is *Ext-injective* in $\text{add } \mathcal{X}$ (the class of finite direct sums of direct summands of modules in $\mathcal{X}$) if $\text{Ext}_\Lambda^1(-, X)|_{\mathcal{X}} = 0$, and *splitting injective* in $\mathcal{X}$ if each injective morphism $X \rightarrow Y$ with Y in $\text{add } \mathcal{X}$ splits. We denote by $I_0(\mathcal{X})$ the class of all indecomposable splitting injectives in $\mathcal{X}$. The following is easy.

Lemma 1. *Let $\mathcal{X}$ be a subcategory of $\text{mod } \Lambda$ closed under extensions and cokernels of injections and let $X \in \mathcal{X}$. Then X is a splitting injective in $\mathcal{X}$ if and only if X is Ext-injective in $\text{add } \mathcal{X}$.*

The following proposition is a particular case of [3, Proposition 3.6].

Proposition 2. *Suppose that $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite. Let $I_1, I_2, \dots, I_n$ be a complete set of non-isomorphic indecomposable injective Λ -modules, and, for each $j = 1, 2, \dots, n$, let $f_j : A_j \rightarrow I_j$ be the unique right minimal $\mathcal{P}_\Lambda^{<\infty}$ -approximations. Then*

$$I_0(\mathcal{P}_\Lambda^{<\infty}) = \cup_{j=1}^n \text{ind } A_j.$$

where $\text{ind } A_j$ denotes the set of indecomposable direct summands of A_j .

Combining this proposition with the lemma, we get:

Corollary 3. *If $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite, then the indecomposable direct summands of the unique right minimal $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the indecomposable injective Λ -modules coincide with the indecomposable Ext-injective modules in $\mathcal{P}_\Lambda^{<\infty}$.*

For a subcategory $\mathcal{X}$ of $\text{mod } \Lambda$, we let

$$\mathcal{X}^\perp = \{M \in \text{Mod } \Lambda \mid \text{Ext}_\Lambda^i(-, M)|_{\mathcal{X}} = 0, \text{ for all } i > 0\}.$$

In particular, the shifting lemma gives

$$(\mathcal{P}_\Lambda^{<\infty})^\perp = \{M \in \text{Mod } \Lambda \mid \text{Ext}_\Lambda^1(-, M)|_{\mathcal{P}_\Lambda^{<\infty}} = 0\}.$$

The following result is the dual of [2, Theorem 5.5 (b)]; see also [5, Theorem 2.1]. Recall that a subcategory $\mathcal{X}$ of $\text{mod } \Lambda$ is said to be *resolving* if it contains the projective Λ -modules and it is closed for extensions and kernels of surjections.

Proposition 4. *The map $\mathcal{X} \mapsto \mathcal{X} \cap \mathcal{X}^\perp$ gives a one-to-one correspondence between contravariantly finite resolving subcategories $\mathcal{X}$ of $\mathcal{P}_\Lambda^{<\infty}$ and equivalence classes of tilting Λ -modules in $\text{mod } \Lambda$.*

In the above proposition, a *tilting module* is an object T in $\text{mod } \Lambda$ such that

- (i) The projective dimension of T is finite;
- (ii) $T \in T^\perp$;
- (iii) there exists a long exact sequence $0 \rightarrow \Lambda_\Lambda \rightarrow T^0 \rightarrow \dots \rightarrow T^n \rightarrow 0$, with $T^i \in \text{add } T$ for all $i = 0, 1, \dots, n$,

and two tilting modules T' and T'' are called *equivalent* if $\text{add } T = \text{add } T'$.

Corollary 5. *If $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite, then the indecomposable Ext-injectives in $\mathcal{P}_\Lambda^{<\infty}$ form a finitely generated tilting Λ -module.*

Proof. Indeed, we have

$$\begin{aligned} (\mathcal{P}_\Lambda^{<\infty}) \cap (\mathcal{P}_\Lambda^{<\infty})^\perp &= \{M \in \mathcal{P}_\Lambda^{<\infty} \mid \text{Ext}_\Lambda^i(-, M)|_{\mathcal{P}_\Lambda^{<\infty}} = 0, \text{ for all } i > 0\} \\ &= \{M \in \mathcal{P}_\Lambda^{<\infty} \mid \text{Ext}_\Lambda^1(-, M)|_{\mathcal{P}_\Lambda^{<\infty}} = 0\} \\ &= \{\text{Ext-injectives in } \mathcal{P}_\Lambda^{<\infty}\}. \end{aligned}$$

□

We also need the following results from [1]. In what follows, $\text{Add } \mathcal{X}$ denotes the class of direct summands of arbitrary direct sums of objects of $\mathcal{X}$.

Lemma 6. [1, Lemma 1.7] *Let T be a tilting module. Then for each M in $T^\perp$ there exists a short exact sequence $0 \longrightarrow K_0 \longrightarrow T_0 \longrightarrow M \longrightarrow 0$, where T_0 is in $\text{Add } T$ and K_0 is in $T^\perp$.*

Theorem 7. [1, Theorems 2.6 and 4.2] *Let Λ be an artin algebra. The following conditions are equivalent:*

- (a) $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite in $\text{mod } \Lambda$;
- (b) there is a finitely presented tilting module T such that $T^\perp = (\mathcal{P}_\Lambda^{<\infty})^\perp$.

Moreover, in this case, the finitistic dimension of Λ is equal to the projective dimension of T .

We can now show our main result.

Theorem 8. *Let Λ be an artin algebra. The following conditions are equivalent:*

- (a) $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite in $\text{mod } \Lambda$;
- (b) the direct sum E of the indecomposable Ext-injective modules in $\mathcal{P}_\Lambda^{<\infty}$ form a tilting Λ -module.

Moreover, in this case, E is the direct sum of the isoclasses of minimal right $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the indecomposable Λ -injectives modules, and the finitistic dimension of Λ is equal to the projective dimension of E .

Proof. First, (a) implies (b) by Corollary 5. Thus, assume that the direct sum E of the Ext-injective modules in $\mathcal{P}_\Lambda^{<\infty}$ forms a tilting Λ -module. We will show that $E^\perp = (\mathcal{P}_\Lambda^{<\infty})^\perp$.

Since $E \in \mathcal{P}_\Lambda^{<\infty}$, we have $(\mathcal{P}_\Lambda^{<\infty})^\perp \subseteq E^\perp$. Conversely, assume that $M \in E^\perp$. We need to show that $\text{Ext}_\Lambda^i(N, M) = 0$ for all $N \in \mathcal{P}_\Lambda^{<\infty}$ and all $i > 0$. So let $N \in \mathcal{P}_\Lambda^{<\infty}$ be of projective dimension n .

Since E is a tilting module and $M \in E^\perp$, it follows from Lemma 6 that there exists a short exact sequence

$$0 \longrightarrow K_0 \longrightarrow E_0 \longrightarrow M \longrightarrow 0,$$

where $E_0 \in \text{Add } E$ and $K_0 \in E^\perp$. Inductively, we get an exact sequence

$$0 \longrightarrow K_{n-1} \longrightarrow E_{n-1} \xrightarrow{f_{n-1}} \cdots \longrightarrow E_1 \xrightarrow{f_1} E_0 \xrightarrow{f_0} M \longrightarrow 0,$$

where $E_i \in \text{Add } E$ and $K_i = \text{Ker } f_i$ lies in $E^\perp$ for all $i > 0$.

By Corollary 3 and [1, Lemma 1.5], each E_i is in $(\mathcal{P}_\Lambda^{<\infty})^\perp$. Therefore, applying the functor $\text{Hom}_\Lambda(N, -)$ to each sequence

$$0 \longrightarrow K_i \longrightarrow E_i \xrightarrow{f_i} K_{i-1} \longrightarrow 0,$$

(where $K_{-1} = M$) yields, for each $j > 0$, an isomorphism

$$\text{Ext}_\Lambda^j(N, K_{i-1}) \cong \text{Ext}_\Lambda^{j+1}(N, K_i).$$

Therefore,

$$\text{Ext}_\Lambda^j(N, M) \cong \text{Ext}_\Lambda^{j+1}(N, K_0) \cong \cdots \cong \text{Ext}_\Lambda^{j+n}(N, K_{n-1}) = 0$$

for all $j > 0$ because $\text{pd } N = n$. Hence $M \in (\mathcal{P}_\Lambda^{<\infty})^\perp$. Therefore $E^\perp = (\mathcal{P}_\Lambda^{<\infty})^\perp$ as claimed, and $\mathcal{P}_\Lambda^{<\infty}$ is contravariantly finite by Theorem 7. Thus (b) implies (a).

Finally, E is the direct sum of the isoclasses of minimal right $\mathcal{P}_\Lambda^{<\infty}$ -approximations of the indecomposable Λ -injectives modules by Corollary 3 and $\text{fin.dim. } \Lambda = \text{pd } E$ by Theorem 7. $\square$

REFERENCES

- [1] L. Angeleri-Hügel and J. Trlifaj, *Tilting theory and the finitistic dimension conjectures*, Trans. Amer. Math. Soc. **354** (2002), no. 11, 4345–4358.
- [2] M. Auslander and I. Reiten, *Applications of contravariantly finite subcategories*, Adv. Math., **86** (1991), no. 1, 111–152.
- [3] M. Auslander and S.O. Smalø, *Preprojective modules over artin algebras*, J. Algebra, **66** (1980), no. 1, 61–122.
- [4] D. Happel and L. Unger, *Modules of finite projective dimension and cocovers*, Math. Ann. **306** (1996), 445–557.
- [5] Ø. Solberg, *Infinite dimensional tilting modules over finite dimensional algebras*, Handbook of tilting theory, 323–344, London Math. Soc. Lecture Note Ser., **332** Cambridge Univ. Press, Cambridge, 2007.

FRANÇOIS HUARD; DEPARTMENT OF MATHEMATICS, BISHOP'S UNIVERSITY. SHERBROOKE, QUÉBEC, CANADA, J1M1Z7.

E-mail address: fhuard@ubishops.ca

DAVID SMITH; DEPARTMENT OF MATHEMATICS, BISHOP'S UNIVERSITY. SHERBROOKE, QUÉBEC, CANADA, J1M1Z7.

E-mail address: dsmith@ubishops.ca